arXiv:1602.04748v2 [math.AT] 23 Jun 2016

BETTI NUMBERS OF UNORDERED CONFIGURATION SPACES OF THE TORUS

CHRISTOPH SCHIESSL

ABSTRACT. Using a method of Félix and Thomas we compute the Betti numbers of unordered configuration spaces of the torus.

1. INTRODUCTION

Let X be a topological space. We denote by

$$F_n(X) = \{(x_1, \dots, x_n) \in X^n \colon x_i \neq x_j \text{ for } i \neq j\}$$

the space of *ordered configurations* of n distinct points on X. The group $\mathfrak{S}_n$ acts freely on $F_n(X)$ by permuting the n points. The quotient

$$C_n(X) = F_n(X)/\mathfrak{S}_n$$

is the space of *unordered configurations* of n points on X.

In the context of representation stability, Church showed that for a connected, orientable manifold M of finite type the rational cohomology groups $H^i(C_n(M), \mathbb{Q})$ stabilise for $n > i$ [Chu12, Cor. 3]. However, very few of these stable Betti numbers have been explicitly computed. Félix and Thomas [FT00] showed that for a closed, orientable, nilpotent, even-dimensional manifold M, the rational Betti numbers of $C_n(M)$ are determined by the rational cohomology algebra $H^*(M, \mathbb{Q})$. They constructed an explicit differential graded algebra that we use to compute the Betti numbers of the unordered configuration spaces of the torus $\Sigma_1 = S^1 \times S^1$.

Theorem. *Suppose $n \geq 2$. Then*

$$\dim_{\mathbb{Q}} H^i(C_n(\Sigma_1), \mathbb{Q}) = \begin{cases} \frac{n-2}{2} & i = n+1,\ n \text{ even} \\ \frac{n+1}{2} & i = n+1,\ n \text{ odd} \\ \frac{3n-4}{2} & i = n,\ n \text{ even} \\ \frac{3n-1}{2} & i = n,\ n \text{ odd} \\ 2i-1 & 2 \leq i < n \\ 2 & i = 1 \\ 1 & i = 0. \end{cases}$$

Date: June 24, 2016.

The author was supported by the grant ERC-2012-AdG-320368-MCSK. Thanks to Gabriel Drummond-Cole, Javier Fresán, Felix Janda, Dan Petersen, Johannes Schmitt, Junliang Shen, Qizheng Yin for very helpful discussions and especially to Rahul Pandharipande for his invaluable support.

Azam [Aza15] determined the rational Betti numbers of configuration spaces of Riemann surfaces for $n = 2, 3$ in any genus and for $n = 4$ in genus 1 by the Kriz model [Kri94]. Napolitano [Nap03] computed the integral cohomology groups of $C_n(\Sigma_1)$ for $n \leq 7$ using a cellular decomposition. Drummond-Cole and Knudsen announced computations of Betti numbers for unordered configuration spaces of surfaces for all genera by a different method [Knu14, p. 30].

The theorem has been tested for all $n \leq 20$ using the computer algebra system SAGE [Dev15].

2. Conventions

We consider $n \geq 2$ as $C_1(X) \simeq X$. We will always work with cohomology/homology with $\mathbb{Q}$-coefficients and identify

$$H^*(M, \mathbb{Q}) = \mathrm{Hom}_{\mathbb{Q}}(H_*(M, \mathbb{Q}), \mathbb{Q})$$

. The free $\mathbb{Q}$-vector space with basis $x_1, \ldots, x_n$ is denoted by $\langle x_1, \ldots, x_n \rangle$.

For any differential graded commutative algebra (A, d), we use the sign convention $xy = (-1)^{\deg x \deg y} yx$ and $d(xy) = d(x)y + (-1)^{\deg x} x d(y)$ for homogenous $x, y \in A$. We have the free graded commutative algebra $\Lambda(V)$ on any graded vector space V with

$$\Lambda(V) = \text{Exterior algebra } (V^{\mathrm{odd}}) \otimes \text{Symmetric algebra } (V^{\mathrm{even}}).$$

3. Construction of the algebra

Let M be a manifold. The cup product gives a map

$$\cup \colon H^*(M) \otimes H^*(M) \to H^*(M),$$

which dualizes to the diagonal comultiplication

$$\Delta \colon H_*(M) \to H_*(M) \otimes H_*(M).$$

Using a basis e_i of $H^*(M)$ the map Δ is given by

$$\Delta(e_k^*) = \sum_{i,j} (\text{coefficient of } e_k \text{ in } e_i \cup e_j)\ e_i^* \otimes e_j^*,$$

where e_i^* denotes the dual basis of $H_*(M)$.

Set $m = \dim(M)$. We take two shifted copies V, W of the vector space $H_*(M)$ with (upper) grading

$$V^{m-r} = H_r(M) \qquad\qquad W^{2m-1-r} = H_r(M).$$

We endow the free graded algebra $\Omega = \Lambda(V \oplus W)$ with the unique differential D of degree 1 such that

$$D_{|V} = 0 \qquad\qquad D_{|W} \colon W \simeq H_*(M) \xrightarrow{\Delta} \Lambda^2 H_*(M) \simeq \Lambda^2 V.$$

A lower grading

$$\Omega = \bigoplus_{n \geq 0} \Omega_n$$

can be defined by putting V in degree 1 and W in degree 2. Hence we have

$$\Omega_n = \bigoplus_{r+2s=n} \Lambda^r V \otimes \Lambda^s W.$$

The vectorspace Ω_n is also graded

$$\Omega_n = \bigoplus_{i \geq 0} \Omega_n^i$$

by the upper grading inherited from Ω. As $D(W) \subset \Lambda^2 V$, the differential D respects the lower grading and Ω_n is a subcomplex of (Ω, D).

Félix and Thomas showed that (Ω_n, D) is a model for the cohomology of $H^*(C_n(M), \mathbb{Q})$.

Theorem 3.1. [FT00, Th. A(2)] *Let M be an orientable, closed, nilpotent, even-dimensional manifold. There is an isomorphism of graded vector spaces*

$$H^*(C_n(M), \mathbb{Q}) \simeq H^*(\Omega_n, D).$$

4. Configuration spaces of the torus

Now we apply this theorem for the torus Σ_1. Its cohomology algebra is $H^*(\Sigma_1) = \langle 1, a, b, ab \rangle$ with $\deg(a) = \deg(b) = 1$ and the relations $ab = -ba$, $a^2 = b^2 = 0$. As $\pi_1(\Sigma_1) = \mathbb{Z}^2$ is abelian and the higher homotopy groups of Σ_1 vanish, Σ_1 is a nilpotent space. We introduce the graded vector spaces $V = \langle v_1, v_a, v_b, v_{ab} \rangle$ and $W = \langle w_1, w_a, w_b, w_{ab} \rangle$ with degrees

$$\begin{aligned} \deg v_1 &= 2 & \deg w_1 &= 3 \\ \deg v_a &= 1 & \deg w_a &= 2 \\ \deg v_b &= 1 & \deg w_b &= 2 \\ \deg v_{ab} &= 0 & \deg w_{ab} &= 1. \end{aligned}$$

We look at the graded algebra $\Omega = \Lambda\langle v_1, v_a, v_b, v_{ab}, w_1, w_a, w_b, w_{ab} \rangle$ with the differential D given by

$$\begin{aligned} D(v_1) &= 0 & D(w_1) &= v_1^2 \\ D(v_a) &= 0 & D(w_a) &= 2v_1 v_a \\ D(v_b) &= 0 & D(w_b) &= 2v_1 v_b \\ D(v_{ab}) &= 0 & D(w_{ab}) &= 2v_1 v_{ab} + 2v_a v_b. \end{aligned}$$

By Theorem 3.1 we have to compute the cohomology groups of the subcomplexes

$$\Omega_n = \bigoplus_{r+2s=n} \Lambda^r V \oplus \Lambda^s W.$$

We will do this by embedding them into the algebra

$$\Theta = \Lambda\langle v_1, v_a, v_b, w_1, w_a, w_b, w_{ab} \rangle,$$

with differential d given by:

$$\begin{aligned} d(v_1) &= 0 & d(w_1) &= v_1^2 \end{aligned}$$

$$\begin{aligned} d(v_a) &= 0 & d(w_a) &= 2v_1v_a \\ d(v_b) &= 0 & d(w_b) &= 2v_1v_b \\ & & d(w_{ab}) &= 2v_1 + 2v_av_b. \end{aligned}$$

All variables have the same grading as in Ω; we only set $v_{ab} = 1$.

Lemma 4.1. *There is an isomorphism $H^i(\Omega_n, D) \simeq H^i(\Theta, d)$ for $i < n$.*

Proof. The injective map

$$\pi\colon \Omega_n \to \Theta, \quad v_{ab} \mapsto 1$$

respects the grading as $\deg v_{ab} = 0$ and commutes with the differentials. Take a degree $i < n$ and any monomial

$$\prod v_k^{e_k} \prod w_l^{f_l} \in \Theta^i$$

of degree i. The only generators of Ω where the lower degree exceeds the upper one are v_{ab} and w_{ab}. As $w_{ab}^2 = 0$ we see

$$\sum e_k + 2\sum f_l \leq i + 1.$$

So the monomial

$$v_{ab}^{n-\sum e_k - 2\sum f_l} \prod v_k^{e_k} \prod w_l^{f_l}$$

is in Ω_n and

$$\pi(v_{ab}^{n-\sum e_k - 2\sum f_l} \prod v_k^{e_k} \prod w_l^{f_l}) = \prod v_k^{e_k} \prod w_l^{f_l}.$$

Thus π is also surjective in degree i. Altogether, π induces an isomorphism $H^i(\Omega_n, d) \simeq H^i(\Theta, d)$ for $i < n$. □

In order to compute the Betti numbers of (Θ, d) we compare d with the simpler differential d_0 given by

$$\begin{aligned} d_0(v_1) &= 0 & d_0(w_1) &= 0 \\ d_0(v_a) &= 0 & d_0(w_a) &= 0 \\ d_0(v_b) &= 0 & d_0(w_b) &= 0 \\ & & d_0(w_{ab}) &= 2v_1 + 2v_av_b \end{aligned}$$

Lemma 4.2. *There is an isomorphism $\varphi\colon (\Theta, d_0) \to (\Theta, d)$.*

Proof. It can be explicitly given by

$$\begin{aligned} \varphi(v_1) &= v_1 & \varphi(w_1) &= w_1 - \frac{1}{2}v_1w_{ab} + \frac{1}{2}v_bw_a \\ \varphi(v_a) &= v_a & \varphi(w_a) &= w_a + v_aw_{ab} \\ \varphi(v_b) &= v_b & \varphi(w_b) &= w_b + v_bw_{ab} \\ & & \varphi(w_{ab}) &= w_{ab} \end{aligned}$$

As $d(\varphi(w_1)) = d(\varphi(w_a)) = d(\varphi(w_b)) = 0$, the map φ commutes with the differentials. □

Lemma 4.3. *The Betti numbers of $H^*(\Theta, d_0)$ are*

$$\dim_{\mathbb{Q}} H^i(\Theta, d_0) = \begin{cases} 1 & i = 0 \\ 2 & i = 1 \\ 2i-1 & i \geq 2. \end{cases}$$

Proof. Denote $T = \Lambda\langle v_1, v_a, v_b, w_1, w_a, w_b\rangle$. Then we have $\Theta = T \oplus Tw_{ab}$. Observe that $d_0|T = 0$. Take homogenous $x, y \in T$. We compute

$$d_0(x + yw_{ab}) = d_0(x) + d_0(y)w_{ab} \pm yd_0(w_{ab}) = \pm 2y(v_1 + v_a v_b).$$

As v_1 has even degree, $v_1 + v_a v_b$ is not a zero-divisor. So we know that $\mathrm{Ker}(d_0) = T$ and

$$H^*(\Theta, d_0) = T/(v_1 + v_a v_b) \simeq T/(v_1) \simeq \Lambda\langle v_a, v_b, w_1, w_a, w_b\rangle.$$

The Poincaré series of $\Lambda\langle v_a, v_b, w_1, w_a, w_b\rangle$ is

$$\frac{(1 + t^{\deg v_a})(1 + t^{\deg v_b})(1 + t^{\deg w_1})}{(1 - t^{\deg w_a})(1 - t^{\deg w_b})} = \frac{(1+t)^2(1+t^3)}{(1-t^2)^2} = \frac{1+t^3}{(1-t)^2},$$

which expands to

$$1 + 2t + 3t^2 + 5t^3 + 7t^4 + \cdots + (2i-1)t^i + \cdots . \qquad \square$$

Combining Lemmas 4.1, 4.2 and 4.3 we have computed $\dim_{\mathbb{Q}} H^i(\Omega_n)$ for $i < n$.

Remark. We consider the morphism

$$p\colon \Omega_n \to \Lambda\langle v_a, v_b, w_1, w_a, w_b, w_{ab}\rangle,\ v_{ab} \mapsto 1,\ v_1 \mapsto -v_a v_b.$$

The above proof shows that for any $x \in \mathrm{Im}\, D$ necessarily $p(x) = 0$.

Lemma 4.4. *We have*

$$\dim_{\mathbb{Q}} H^{n+1}(\Omega_n) = \begin{cases} \frac{n-2}{2} & n \text{ even} \\ \frac{n+1}{2} & n \text{ odd} \end{cases} \qquad \dim_{\mathbb{Q}} H^i(\Omega_n) = 0 \text{ for } i > n+1.$$

Proof. We denote $\Theta' = \Lambda\langle v_1, v_a, v_b, v_{ab}, w_a, w_b, w_{ab}\rangle$. The only generators with upper grading exceeding the lower grading are v_1 and w_1. Hence any $x \in \Omega_n^i$ with $i > n$ can be written as $x = v_1 f + w_1 g$ where $f, g \in \Theta'$. We compute

$$D(x) = v_1 D(f) + v_1^2 g - w_1 D(g).$$

As $D(\Theta') \subset \Theta'$ we see that $D(x) = 0$ implies $D(g) = 0$. So $x \in \mathrm{Ker}\, D$ if and only if $D(f) = -v_1 g$. Therefore any $x \in \mathrm{Ker}\, D$ is of the form

$$x(f) = v_1 f - w_1 \frac{D(f)}{v_1}$$

with $f \in \Theta'$ such that $v_1 | D(f)$.

We will now discuss when the cycles $x(f)$ are a boundary. If $f = v_1 h$ then

$$D(w_1 h) = v_1^2 h - w_1 D(h) = v_1 f - w_1 \frac{D(f)}{v_1} = x(f).$$

For any $x(f) \in \Omega^i_n$ with $i > n+1$ we know that f has to be divisible by v_1. Hence $H^i(\Omega_n) = 0$ for $i > n+1$.

Now we look at the case $i = n+1$. If $v_{ab}|f$ or $w_{ab}|f$ then $v_1|f$ for degree reasons.

We consider the sets

$$B_{\text{odd}} = \{w_a^{n_1} w_b^{n_2} \mid 2n_1 + 2n_2 + 1 = n;\, n_1, n_2 \geq 0\}$$

for odd degree n and

$$B_{\text{even}} = \{v_b w_a w_a^{n_1} w_b^{n_2} \mid 2n_1 + 2n_2 + 4 = n;\, n_1, n_2 \geq 0\}$$

for even n.

If $v_1 f = D(h)$ then $D(f) = 0$ and hence we get $x(f) = D(h) \in \mathrm{Im}(D)$. Using the relations

$$\begin{aligned} D(v_b w_a^{n_1+1} w_b^{n_2}) &= -2(n_1+1) v_1 v_a v_b w_a^{n_1} w_b^{n_2} \\ D(w_a^{n_1+1}) &= 2(n_1+1) v_1 v_a w_a^{n_1} \\ D(w_b^{n_1+1}) &= 2(n_1+1) v_1 v_b w_b^{n_1} \\ D(w_a^{n_1+1} w_b^{n_2+1}) &= 2(n_1+1) v_1 v_a w_a^{n_1} w_b^{n_2+1} + 2(n_2+1) v_1 v_b w_a^{n_1+1} w_b^{n_2} \end{aligned}$$

we conclude that the set $\{x(b) | b \in B_{\text{even}}\}$ resp. $\{x(b) | b \in B_{\text{odd}}\}$ is a generating system of $H^{n+1}(\Omega_n)$ for even resp. odd n.

By applying p we see that no non-trivial linear combinations of these generating sets are boundaries. Hence we found an explicit basis of $H^{n+1}(\Omega_n)$. □

Lemma 4.5. *We have*

$$\dim_{\mathbb{Q}} H^n(\Omega_n) = \begin{cases} \frac{3n-4}{2} & n \text{ even} \\ \frac{3n-1}{2} & n \text{ odd} \end{cases}.$$

Proof. As the torus acts freely on $C_n(\Sigma_1)$ we have $\chi(\Omega_n) = 0$. Using the above computation of $\dim_{\mathbb{Q}} H^{n+1}(\Omega_n)$ and

$$\sum_{i=0}^{n-1} \dim_{\mathbb{Q}} H^i(\Omega_n) = 1 - 2 + 3 + \cdots + (-1)^{n-1}(2n-3) = (-1)^{n-1}(n-1)$$

we can reconstruct the only missing Betti number $\dim_{\mathbb{Q}} H^n(\Omega_n)$. □

Combining all lemmas, we have computed $\dim_{\mathbb{Q}} H^i(C_n(\Sigma_1), \mathbb{Q})$ for all i. We reproduce exactly the stability result

$$\dim_{\mathbb{Q}} H^i(C_{n+1}(\Sigma_1), \mathbb{Q}) = \dim_{\mathbb{Q}} H^i(C_n(\Sigma_1), \mathbb{Q})$$

for $n > i$ of Church [Chu12, Cor. 3].

Remark. Let $d \geq 1$. With the same method one immediately finds for $n \geq 3$

$$\dim_{\mathbb{Q}} H^i(C_n(S^{2d}), \mathbb{Q}) = \begin{cases} 1 & \text{for } i = 0,\, 4d-1 \\ 0 & \text{otherwise,} \end{cases}$$

which has also been computed by [RW13], [Sal04].

Remark. It seems that our method does not work for surfaces of genus $g > 1$ because the differential can not be deformed as in Lemma 4.2.

References

[Aza15] Haniya Azam. Cohomology groups of configuration spaces of Riemann surfaces. *Bull. Math. Soc. Sci. Math. Roumanie (N.S.)*, 58(106)(1):33–47, 2015.

[Chu12] Thomas Church. Homological stability for configuration spaces of manifolds. *Invent. Math.*, 188(2):465–504, 2012.

[Dev15] The Sage Developers. *Sage Mathematics Software (Version 6.7)*, 2015. `http://www.sagemath.org`.

[FT00] Yves Félix and Jean-Claude Thomas. Rational Betti numbers of configuration spaces. *Topology Appl.*, 102(2):139–149, 2000.

[Knu14] Ben Knudsen. Betti numbers and stability for configuration spaces via factorization homology. *ArXiv e-prints*, May 2014.

[Kri94] Igor Kriz. On the rational homotopy type of configuration spaces. *Ann. of Math.*, 139(2):227–237, 1994.

[Nap03] Fabien Napolitano. On the cohomology of configuration spaces on surfaces. *J. Lond. Math. Soc.*, 68(2):477–492, 2003.

[RW13] Oscar Randal-Williams. Topological chiral homology and configuration spaces of spheres. *Morfismos*, 17(2):57–69, 2013.

[Sal04] Paolo Salvatore. Configuration spaces on the sphere and higher loop spaces. *Math. Z.*, 248(3):527–540, 2004.

Departement Mathematik, ETH Zürich, Rämistrasse 101, 8092 Zürich, Switzerland

E-mail address: `christoph.schiessl@math.ethz.ch`